SPECIFICATION

Applicant herein requests that the following amendments be made to page 1 and page 6-7 of the Specification of this application. Applicant respectfully requests that current paragraph 1 of page 1 be substituted for the following amended paragraph and that the current incomplete last paragraph of page 6 and first incomplete paragraph of page 7 be substituted with the following amended paragraph. No new matter is added by these amendments.

Page 1, Paragraph 1: marked-up version

Field of Invention

The present invention relates to a novel process for decolorization of colored effluents. More particularly it relates to a process for decolorization of colored effluents of textile mills, dye-making industries, paper and pulp industries and molasses spent wash from alcohol distilleries using an unidentified white-rot marine fungus NIOCC #2a isolated from mangrove wood and deposited on September 7, 2004 in the microbial type culture collection (MTCC) of the Institute of Microbial Technology, Chandigarh, India, under the accession number MTCC 5159. Further, this invention relates to decolorization of these effluents using the fungus directly, its cell-free culture supernatant or immobilized fungus or extracellular polymeric substances produced by the fungus. Furthermore, the decolorization of effluents can be carried out from 30°C to 60°C and at pH 3 to 6. The decolorization of various colored effluents occurs in the presence of sea water with 25 parts per thousand salinity. Several synthetic dyes are also decolorized under similar conditions of temperature and pH by using free mycelia or immobilized fungus or extracellular culture fluids or extracellular polymeric substances.

Clean Version

Field of Invention

The present invention relates to a novel process for decolorization of colored effluents. More particularly it relates to a process for decolorization of colored effluents of textile mills, dye-making industries, paper and pulp industries and molasses spent wash from alcohol distilleries using an unidentified white-rot marine fungus NIOCC #2a isolated from mangrove wood and deposited on September 7, 2004 in the microbial type culture collection (MTCC) of the Institute of Microbial Technology, Chandigarh, India, under the accession number MTCC 5159. Further, this invention relates to decolorization of these effluents using the fungus directly, its cell-free culture supernatant or immobilized fungus or extracellular polymeric substances produced by the fungus. Furthermore, the decolorization of effluents can be carried out from 30°C to 60°C and at pH 3 to 6. The decolorization of various colored effluents occurs in the presence of sea water with 25 parts per thousand salinity. Several synthetic dyes are also decolorized under similar conditions of temperature and pH by using free mycelia or immobilized fungus or extracellular culture fluids or extracellular polymeric substances.

大腿,在大腿的心态的大路下上,在这个人的大腿,那样有有一点的大腿的大腿,一下一下一下,这个人的一下一下,这个人的一个人的一个人的一个人的一个人。 "我们,我不是你们的一个人的人,我们就是一个人的人的,我们的人,我们就是一个人的人的人的人的人的人,我们们也不是一个人的人的人的人,也是一个人的人们也是一个人的

the control of the second of the first of the control of the contr

Page 6 last incomplete paragraph and page 7. first incomplete paragraph

Marked-Up Version:

The most efficient micro organisms to break down colored pollutants so far reported are whiterot fungi. These comprise mostly basidiomycetous fungi capable of extensive aerobic lignin
degradation and mineralization. This is possible through several extra cellular lignin-degrading
enzymes synthesized by these fungi. These enzymes lack substrate specificity and are thus
capable of degrading a wide range of xenobiotics. Thus this invention relates to degradation of
several such xenobiotic compounds in colored effluents by the unidentified white-rot fungus

Serial No. 10/570,749

Page 9 of 16

Attorney Docket No. 007292-01 US

NIOCC #2a deposited at the Institute of Microbial Technology, Chandigarh, India and having the accession No. MTCC 5159 can be made available to public after observing normal established procedures.

Clean Version:

The most efficient micro organisms to break down colored pollutants so far reported are whiterot fungi. These comprise mostly basidiomycetous fungi capable of extensive aerobic lignin
degradation and mineralization. This is possible through several extra cellular lignin-degrading
enzymes synthesized by these fungi. These enzymes lack substrate specificity and are thus
capable of degrading a wide range of xenobiotics. Thus this invention relates to degradation of
several such xenobiotic compounds in colored effluents by the unidentified white-rot fungus.

NIOCC #2a deposited at the Institute of Microbial Technology, Chandigarh, India and having
the accession No. MTCC 5159 can be made available to public after observing normal
established procedures.

REMARKS

Serial No. 10/570,749

Page 10 of 16

Attorney Docket No. 007292-01 US